



Guidelines for the Use of Antiretroviral Agents in Adults and Adolescents with HIV

Downloaded from <https://aidsinfo.nih.gov/guidelines> on 8/31/2020

Visit the AIDSinfo website to access the most up-to-date guideline.

Register for e-mail notification of guideline updates at <https://aidsinfo.nih.gov/e-news>.

Table 22a. Interactions Between Non-Nucleoside Reverse Transcriptase Inhibitors and Protease Inhibitors (Last updated December 18, 2019; last reviewed December 18, 2019) (page 1 of 2)

Note: Interactions associated with DLV, FPV, IDV, NFV, and SQV are **not** included in this table. Please refer to the FDA product labels for information regarding interactions between these drugs and other concomitant drugs.

PIs		NNRTIs				
		DOR	EFV	ETR	NVP	RPV
ATV Unboosted	PK Data	↑ DOR expected ↔ ATV expected	↔ EFV ATV AUC ↓ 74%	ETR AUC ↑ 50% and C _{min} ↑ 58% ↔ ATV AUC and C _{min} ↓ 47%	↑ NVP possible ↓ ATV possible	↑ RPV possible ↔ ATV expected
	Dose	No dose adjustment needed.	Do not coadminister.	Do not coadminister.	Do not coadminister.	No dose adjustment needed.
ATV/c	PK Data	↑ DOR expected ↔ ATV expected	↔ EFV expected ↓ ATV possible ↓ COBI possible	↑ ETR possible ↓ ATV possible ↓ COBI possible	↑ NVP possible ↓ ATV possible ↓ COBI possible	↑ RPV possible ↔ ATV expected
	Dose	No dose adjustment needed.	ATV/c in ART-Naive Patients: <ul style="list-style-type: none"> • ATV 400 mg plus COBI 150 mg once daily • Do not use coformulated ATV 300 mg/COBI 150 mg. ATV/c in ART-Experienced Patients: <ul style="list-style-type: none"> • Do not coadminister. No dose adjustment needed for EFV.	Do not coadminister.	Do not coadminister.	No dose adjustment needed.
ATV/r	PK Data	↑ DOR expected ↔ ATV expected	↔ EFV expected (ATV 400 mg plus RTV 100 mg) Once Daily: <ul style="list-style-type: none"> • ATV concentrations similar to (ATV 300 mg plus RTV 100 mg) without EFV 	(ATV 300 mg plus RTV 100 mg) Once Daily: <ul style="list-style-type: none"> • ETR AUC and C_{min} both ↑ ~30% • ↔ ATV AUC and C_{min} 	(ATV 300 mg plus RTV 100 mg) Once Daily: <ul style="list-style-type: none"> • ATV AUC ↓ 42% and C_{min} ↓ 72% • NVP AUC ↑ 25% 	↑ RPV possible ↔ ATV expected
	Dose	No dose adjustment needed.	ATV/r in ART-Naive Patients: <ul style="list-style-type: none"> • (ATV 400 mg plus RTV 100 mg) once daily ATV/r in ART-Experienced Patients: <ul style="list-style-type: none"> • Do not coadminister. No dose adjustment needed for EFV.	No dose adjustment needed.	Do not coadminister.	No dose adjustment needed.
DRV/c	PK Data	↑ DOR expected ↔ DRV expected	↔ EFV expected ↓ DRV possible ↓ COBI possible	ETR 400 mg Once Daily with (DRV 800 mg plus COBI 150 mg) Once Daily: <ul style="list-style-type: none"> • ↔ ETR AUC and C_{min} • ↔ DRV AUC and C_{min} ↓ 56% • COBI AUC ↓ 30% and C_{min} ↓ 66% 	↑ NVP possible ↓ DRV possible ↓ COBI possible	↔ DRV expected ↑ RPV possible
	Dose	No dose adjustment needed.	Do not coadminister.	Do not coadminister.	Do not coadminister.	No dose adjustment needed.

Table 22a. Interactions Between Non-Nucleoside Reverse Transcriptase Inhibitors and Protease Inhibitors (Last updated December 18, 2019; last reviewed December 18, 2019) (page 2 of 2)

PIs		NNRTIs				
		DOR	EFV	ETR	NVP	RPV
DRV/r	PK Data	↑ DOR expected ↔ DRV expected	With (DRV 300 mg plus RTV 100 mg) Twice Daily: • EFV AUC ↑ 21% • ↔ DRV AUC and C _{min} ↓ 31%	ETR 100 mg Twice Daily with (DRV 600 mg plus RTV 100 mg) Twice Daily: • ETR AUC ↓ 37% and C _{min} ↓ 49% • ↔ DRV	With (DRV 400 mg plus RTV 100 mg) Twice Daily: • NVP AUC ↑ 27% and C _{min} ↑ 47% • DRV AUC ↑ 24% ^a	RPV 150 mg Once Daily with (DRV 800 mg plus RTV 100 mg) Once Daily: • RPV AUC ↑ 130% and C _{min} ↑ 178% • ↔ DRV
	Dose	No dose adjustment needed.	Clinical significance unknown. Use standard doses and monitor patient closely. Consider monitoring drug levels.	No dose adjustment needed. Despite reduced ETR concentration, safety and efficacy of this combination have been established in a clinical trial.	No dose adjustment needed.	No dose adjustment needed.
LPV/r	PK Data	↑ DOR expected ↔ LPV expected	↔ EFV expected With LPV/r 500 mg/125 mg ^b Twice Daily: • LPV concentration similar to that of LPV/r 400 mg/100 mg twice daily without EFV	ETR AUC ↓ 35% (comparable to the decrease seen with DRV/r) ↔ LPV AUC	↑ NVP possible LPV AUC ↓ 27% and C _{min} ↓ 51%	RPV 150 mg Once Daily with LPV/r: • RPV AUC ↑ 52% and C _{min} ↑ 74% • ↔ LPV
	Dose	No dose adjustment needed.	LPV/r 500 mg/125 mg ^b twice daily LPV/r 533 mg/133 mg twice daily when using oral solution No dose adjustment needed for EFV.	No dose adjustment needed.	LPV/r 500 mg/125 mg ^b twice daily LPV/r 533 mg/133 mg twice daily when using oral solution No dose adjustment needed for NVP.	No dose adjustment needed.
TPV/r Note: Always use TPV with RTV	PK Data	↑ DOR expected ↔ TPV expected	With (TPV 500 mg plus RTV 100 mg) Twice Daily: • ↔ EFV • TPV AUC ↓ 31% and C _{min} ↓ 42% With (TPV 750 mg plus RTV 200 mg) Twice Daily: • ↔ EFV and TPV	With (TPV 500 mg plus RTV 200 mg) Twice Daily: • ETR AUC ↓ 76% and C _{min} ↓ 82% • ↔ TPV AUC and C _{min} ↑ 24%	With (TPV 250 mg plus RTV 200 mg) Twice Daily or with (TPV 750 mg plus RTV 100 mg) Twice Daily: • ↔ NVP • ↔ TPV expected	↑ RPV possible ↔ TPV expected
	Dose	No dose adjustment needed.	No dose adjustment needed.	Do not coadminister.	No dose adjustment needed.	No dose adjustment needed.

^a DRV concentration was compared to a historic control.

^b Use a combination of two LPV/r 200 mg/50 mg tablets plus one LPV/r 100 mg/25 mg tablet to make a total dose of LPV/r 500 mg/125 mg.

Key to Symbols: ↑ = increase ↓ = decrease ↔ = no change

Key: ART = antiretroviral therapy; ATV = atazanavir; ATV/c = atazanavir/cobicistat; ATV/r = atazanavir/ritonavir; AUC = area under the curve; C_{min} = minimum plasma concentration; COBI = cobicistat; DLV = delavirdine; DOR = doravirine; DRV = darunavir; DRV/c = darunavir/cobicistat; DRV/r = darunavir/ritonavir; EFV = efavirenz; ETR = etravirine; FDA = Food and Drug Administration; FPV = fosamprenavir; IDV = indinavir; LPV = lopinavir; LPV/r = lopinavir/ritonavir; NFV = nelfinavir; NNRTI = non-nucleoside reverse transcriptase inhibitor; NVP = nevirapine; PI = protease inhibitor; PK = pharmacokinetic; RPV = rilpivirine; RTV = ritonavir; SQV = saquinavir; TPV = tipranavir; TPV/r = tipranavir/ritonavir